Leader in Biomolecular Solutions for Life Science

# FOS Rabbit pAb

Catalog No.: A17351 1 Publications



### **Basic Information**

# **Observed MW**

62kDa

#### **Calculated MW**

41kDa

#### Category

Polyclonal Antibody

### **Applications**

WB,ELISA

#### **Cross-Reactivity**

Human, Mouse, Rat

# **Background**

The Fos gene family consists of 4 members: FOS, FOSB, FOSL1, and FOSL2. These genes encode leucine zipper proteins that can dimerize with proteins of the JUN family, thereby forming the transcription factor complex AP-1. As such, the FOS proteins have been implicated as regulators of cell proliferation, differentiation, and transformation. In some cases, expression of the FOS gene has also been associated with apoptotic cell death.

# **Recommended Dilutions**

WB

1:500 - 1:1000

# **Immunogen Information**

Gene ID 2353 Swiss Prot P01100

#### **Immunogen**

A synthetic peptide corresponding to a sequence within amino acids 1-100 of human FOS (NP 005243.1).

### **Synonyms**

p55; AP-1; C-FOS; FOS

# **Contact**

•

www.abclonal.com

# **Product Information**


**Source** Rabbit **Isotype** IgG **Purification**Affinity purification

# Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

# **Validation Data**



Western blot analysis of extracts of various cell lines, using FOS antibody (A17351) at 1:1000 dilution.NIH/3T3 cells were treated by Serum-starvation overnight at  $37^{\circ}$ C.NIH/3T3 cells were treated by PMA/TPA (200 nM) at  $37^{\circ}$ C for 15 minutes after serum-starvation overnight.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 30s.