Leader in Biomolecular Solutions for Life Science

ATRX Rabbit pAb

Catalog No.: A17378



Basic Information

Observed MW

310kDa

Calculated MW

283kDa

Category

Polyclonal Antibody

Applications

WB, ELISA

Cross-Reactivity

Human, Mouse

Background

The protein encoded by this gene contains an ATPase/helicase domain, and thus it belongs to the SWI/SNF family of chromatin remodeling proteins. This protein is found to undergo cell cycle-dependent phosphorylation, which regulates its nuclear matrix and chromatin association, and suggests its involvement in the gene regulation at interphase and chromosomal segregation in mitosis. Mutations in this gene are associated with X-linked syndromes exhibiting cognitive disabilities as well as alpha-thalassemia (ATRX) syndrome. These mutations have been shown to cause diverse changes in the pattern of DNA methylation, which may provide a link between chromatin remodeling, DNA methylation, and gene expression in developmental processes. Multiple alternatively spliced transcript variants encoding distinct isoforms have been reported.

Recommended Dilutions

WB

1:2000 - 1:6000

Immunogen Information

Gene ID

546

Swiss Prot P46100

Immunogen

A synthetic peptide corresponding to a sequence within amino acids 2100-2200 of human ATRX (NP 000480.3).

Synonyms

JMS; XH2; XNP; MRX52; RAD54; RAD54L; ZNF-HX; ATRX

Contact

•

www.abclonal.com


Product Information

Source Rabbit **Isotype** IgG **Purification**Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.



Western blot analysis of various lysates, using ATRX Rabbit pAb (A17378) at 1:5000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (RM00021).

Negative control (NC): U-2 OS.

Exposure time: 60s.

Western blot analysis of Mouse brain, using ATRX Rabbit pAb (A17378) at 1:5000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST. Detektion: Land Eathanced Kit (RM00021).

