Leader in Biomolecular Solutions for Life Science

# MRPL34 Rabbit pAb

Catalog No.: A17761



# **Basic Information**

### **Observed MW**

10kDa

### **Calculated MW**

10kDa

#### Category

Polyclonal Antibody

#### **Applications**

WB,IF/ICC,ELISA

#### **Cross-Reactivity**

Human, Mouse, Rat

# **Background**

Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes and help in protein synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalian mitoribosomes and prokaryotic ribosomes is that the latter contain a 5S rRNA. Among different species, the proteins comprising the mitoribosome differ greatly in sequence, and sometimes in biochemical properties, which prevents easy recognition by sequence homology. This gene encodes a 39S subunit protein.

# **Recommended Dilutions**

**WB** 1:500 - 1:2000

IF/ICC 1:50 - 1:200

# **Immunogen Information**

**Gene ID**Swiss Prot
64981
Q9BQ48

#### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 16-92 of human MRPL34 (NP\_076426.1).

### **Synonyms**

L34mt; MRPL34

### **Contact**

www.abclonal.com

# **Product Information**


SourceIsotypePurificationRabbitIgGAffinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

# **Validation Data**




Western blot analysis of various lysates using MRPL34 Rabbit pAb (A17761) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.



Immunofluorescence analysis of L929 cells using MRPL34 Rabbit pAb (A17761) at dilution of 1:100. Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.