

A18022

Leader in Biomolecular Solutions for Life Science



[KO Validated] Peroxiredoxin 3 (PRDX3) Rabbit pAb

Catalog No.: A18022 **KO** Validated

Basic Information

Observed MW

25kDa

Calculated MW

28kDa

Category

Polyclonal Antibody

Applications

WB, ELISA

Cross-Reactivity

Human, Mouse

Background

This gene encodes a mitochondrial protein with antioxidant function. The protein is similar to the C22 subunit of *Salmonella typhimurium* alkylhydroperoxide reductase, and it can rescue bacterial resistance to alkylhydroperoxide in *E. coli* that lack the C22 subunit. The human and mouse genes are highly conserved, and they map to the regions syntenic between mouse and human chromosomes. Sequence comparisons with recently cloned mammalian homologs suggest that these genes consist of a family that is responsible for the regulation of cellular proliferation, differentiation and antioxidant functions. This family member can protect cells from oxidative stress, and it can promote cell survival in prostate cancer. Alternative splicing of this gene results in multiple transcript variants. Related pseudogenes have been identified on chromosomes 1, 3, 13 and 22.

Recommended Dilutions

WB 1:500 - 1:2000

Immunogen Information

Gene ID

10935

Swiss Prot

P30048

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 63-256 of human Peroxiredoxin 3 (Peroxiredoxin 3 (PRDX3)) (NP_006784.1).

Synonyms

AOP1; MER5; AOP-1; PPPCD; SP-22; HBC189; SCAR32; PRO1748; prx-III; 3)

Contact



www.abclonal.com

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal, 50% glycerol, pH 7.3.

Validation Data



Western blot analysis of lysates from wild type (WT) and Peroxiredoxin 3 (PRDX3) knockout (KO) HeLa cells, using [KO Validated] Peroxiredoxin 3 (PRDX3) Rabbit pAb (A18022) at 1:3000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25 μ g per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 1min.