A18038

Leader in Biomolecular Solutions for Life Science

# [KO Validated] Cortactin Rabbit pAb

Catalog No.: A18038 KO Validated



#### **Basic Information**

Observed MW 80-85kDa

Calculated MW 62kDa

Category Polyclonal Antibody

Applications WB,ELISA

Cross-Reactivity Human,Mouse,Rat

#### Background

This gene is overexpressed in breast cancer and squamous cell carcinomas of the head and neck. The encoded protein is localized in the cytoplasm and in areas of the cell-substratum contacts. This gene has two roles: (1) regulating the interactions between components of adherens-type junctions and (2) organizing the cytoskeleton and cell adhesion structures of epithelia and carcinoma cells. During apoptosis, the encoded protein is degraded in a caspase-dependent manner. The aberrant regulation of this gene contributes to tumor cell invasion and metastasis. Three splice variants that encode different isoforms have been identified for this gene.

#### **Recommended Dilutions**

#### Immunogen Information

WB

1:500 - 1:2000

**Gene ID** 2017

Swiss Prot Q14247

#### Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 1-200 of human Cortactin (NP\_612632.1).

Synonyms EMS1; in

Contact

€

onal.com Source

## Product Information

www.abclonal.com


**lsotype** IgG Purification Affinity purification

Storage

Rabbit

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

### Validation Data



Western blot analysis of lysates from wild type (WT) and Cortactin knockout (KO) HeLa cells, using [KO Validated] Cortactin Rabbit pAb (A18038) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 5s.