#### A18044

Leader in Biomolecular Solutions for Life Science

# [KO Validated] FADD Rabbit pAb

Catalog No.: A18044

KO Validated 1 Publications

## **Basic Information**

**Observed MW** 28kDa

**Calculated MW** 23kDa

Category Polyclonal Antibody

Applications WB, ELISA

**Cross-Reactivity** Human, Mouse, Rat

#### Background

The protein encoded by this gene is an adaptor molecule that interacts with various cell surface receptors and mediates cell apoptotic signals. Through its C-terminal death domain, this protein can be recruited by TNFRSF6/Fas-receptor, tumor necrosis factor receptor, TNFRSF25, and TNFSF10/TRAIL-receptor, and thus it participates in the death signaling initiated by these receptors. Interaction of this protein with the receptors unmasks the Nterminal effector domain of this protein, which allows it to recruit caspase-8, and thereby activate the cysteine protease cascade. Knockout studies in mice also suggest the importance of this protein in early T cell development.

### **Recommended Dilutions**

#### Immunogen Information

WB

1:500 - 1:2000

Gene ID 8772

**Swiss Prot** 013158

#### Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 1-208 of human FADD (NP 003815.1).

Synonyms GIG3; IMD90; MORT1; DD

#### Contact

## **Product Information**

€

www.abclonal.com

Isotype lgG

Purification Affinity purification

Storage


Source

Rabbit

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thimerosal, 50% glycerol, pH7.3.



## Validation Data



Western blot analysis of lysates from wild type (WT) and FADD knockout (KO) HeLa cells, using [KO Validated] FADD Rabbit pAb (A18044) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 5s.