Leader in Biomolecular Solutions for Life Science

HDAC1 Mouse mAb

Catalog No.: A18304 2 Publications



Basic Information

Observed MW

62kDa

Calculated MW

Category

Mouse Monoclonal Antibody

Applications

WB, ELISA

Cross-Reactivity

Human, Mouse

CloneNo number

AMC0232

Background

Histone acetylation and deacetylation, catalyzed by multisubunit complexes, play a key role in the regulation of eukaryotic gene expression. The protein encoded by this gene belongs to the histone deacetylase/acuc/apha family and is a component of the histone deacetylase complex. It also interacts with retinoblastoma tumor-suppressor protein and this complex is a key element in the control of cell proliferation and differentiation. Together with metastasis-associated protein-2, it deacetylates p53 and modulates its effect on cell growth and apoptosis.

Recommended Dilutions

WB

1:500 - 1:2000

Immunogen Information

Gene ID

Swiss Prot

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 393-482 of human HDAC1 (Q13547).

Synonyms

Contact

•

www.abclonal.com

Product Information


Source Mouse **Isotype** IgG1,Kappa **Purification**Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

Validation Data



Western blot analysis of various lysates using HDAC1 Mouse mAb (A18304) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Mouse IgG (H+L) (AS003) at 1:10000 dilution.

Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 30s.