

A18323

Leader in Biomolecular Solutions for Life Science



# JAK1 Rabbit pAb

Catalog No.: A18323

15 Publications

## Basic Information

### Observed MW

133kDa

### Calculated MW

133kDa

### Category

Mouse Monoclonal Antibody

### Applications

WB,IF/ICC,ELISA

### Cross-Reactivity

Human,Mouse,Rat

## Background

This gene encodes a membrane protein that is a member of a class of protein-tyrosine kinases (PTK) characterized by the presence of a second phosphotransferase-related domain immediately N-terminal to the PTK domain. The encoded kinase phosphorylates STAT proteins (signal transducers and activators of transcription) and plays a key role in interferon-alpha/beta, interferon-gamma, and cytokine signal transduction. This gene plays a crucial role in effecting the expression of genes that mediate inflammation, epithelial remodeling, and metastatic cancer progression. This gene is a key component of the interleukin-6 (IL-6)/JAK1/STAT3 immune and inflammation response and is a therapeutic target for alleviating cytokine storms. The kinase activity of this gene is directly inhibited by the suppressor of cytokine signalling 1 (SOCS1) protein. Alternative splicing results in multiple transcript variants.

## Recommended Dilutions

WB 1:1000 - 1:5000

IF/ICC 1:100 - 1:500

## Immunogen Information

### Gene ID

3716

### Swiss Prot

P23458

### Immunogen

A synthetic peptide corresponding to a sequence within amino acids 30-300 of human JAK1 (NP\_002218.2).

### Synonyms

JTK3; AIIDE; JAK1A; JAK1B; JAK1

## Contact



[www.abclonal.com](http://www.abclonal.com)

## Product Information

### Source

Rabbit

### Isotype

IgG

### Purification

Affinity purification


### Storage

Store at -20°C. Avoid freeze / thaw cycles.


Buffer: PBS with 0.01% thimerosal, 50% glycerol, pH7.3.

## Validation Data

---



Western blot analysis of NIH/3T3, using JAK1 Rabbit pAb (A18323) at 1:2000 dilution.  
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (A5014) at 1:10000 dilution.  
Lysates/proteins: 25µg per lane.  
Blocking buffer: 3% nonfat dry milk in TBST.  
Detection: ECL Basic Kit (RM00020).  
Exposure time: 90s.



Immunofluorescence analysis of U2OS cells using JAK1 Rabbit pAb (A18323) at dilution of 1:300 (40x lens). Blue: DAPI for nuclear staining.