Leader in Biomolecular Solutions for Life Science

# GC1q R/C1QBP Rabbit pAb

Catalog No.: A1883 3 Publications



### **Basic Information**

# **Observed MW**

36kDa

### **Calculated MW**

31kDa

#### Category

Polyclonal Antibody

#### **Applications**

WB,IF/ICC,ELISA

#### **Cross-Reactivity**

Human, Mouse, Rat

# **Background**

The human complement subcomponent C1q associates with C1r and C1s in order to yield the first component of the serum complement system. The protein encoded by this gene is known to bind to the globular heads of C1q molecules and inhibit C1 activation. This protein has also been identified as the p32 subunit of pre-mRNA splicing factor SF2, as well as a hyaluronic acid-binding protein.

# **Recommended Dilutions**

**WB** 1:500 - 1:2000

**IF/ICC** 1:50 - 1:200

# **Immunogen Information**

**Gene ID**708

Swiss Prot
Q07021

#### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 74-282 of human GC1q R/GC1q R/C1QBP (NP\_001203.1).

### **Synonyms**

p32; HABP1; gC1qR; GC1QBP; SF2p32; gC1Q-R; COXPD33; SF2AP32; GC1q R/C1QBP

### **Contact**

www.abclonal.com

# **Product Information**


SourceIsotypePurificationRabbitIgGAffinity purification

# Storage

Store at -20°C. Avoid freeze / thaw cycles.


Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

# **Validation Data**



Western blot analysis of various lysates using GC1q R/C1QBP Rabbit pAb (A1883) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit lgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins:  $25\mu g$  per lane.

Blocking buffer: 3% nonfat dry milk in TBST.



Immunofluorescence analysis of HeLa cells using GC1q R/C1QBP Rabbit pAb (A1883). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.