

## Doublecortin Rabbit mAb

Catalog No.: A19042    Recombinant    1 Publications

### Basic Information

**Observed MW**

45kDa

**Calculated MW**

41kDa

**Category**

SMab Recombinant Monoclonal Antibody

**Applications**

WB,ELISA

**Cross-Reactivity**

Mouse,Rat

**CloneNo** number

ARC0380

### Background

This gene encodes a member of the doublecortin family. The protein encoded by this gene is a cytoplasmic protein and contains two doublecortin domains, which bind microtubules. In the developing cortex, cortical neurons must migrate over long distances to reach the site of their final differentiation. The encoded protein appears to direct neuronal migration by regulating the organization and stability of microtubules. In addition, the encoded protein interacts with LIS1, the regulatory gamma subunit of platelet activating factor acetylhydrolase, and this interaction is important to proper microtubule function in the developing cortex. Mutations in this gene cause abnormal migration of neurons during development and disrupt the layering of the cortex, leading to epilepsy, cognitive disability, subcortical band heterotopia ("double cortex" syndrome) in females and lissencephaly ("smooth brain" syndrome) in males. Multiple transcript variants encoding different isoforms have been found for this gene.

### Recommended Dilutions

WB                          1:500 - 1:1000

### Immunogen Information

**Gene ID**

1641

**Swiss Prot**

O43602

**Immunogen**

A synthetic peptide corresponding to a sequence within amino acids 1-100 of human Doublecortin (O43602).

**Synonyms**

DC; DBCN; LISX; SCLH; XLIS; Doublecortin

### Contact



[www.abclonal.com](http://www.abclonal.com)

### Product Information

**Source**

Rabbit

**Isotype**

IgG

**Purification**


Affinity purification

**Storage**

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 0.05% BSA, 50% glycerol, pH7.3.

## Validation Data



Western blot analysis of extracts of various cell lines, using Doublecortin antibody (A19042) at 1:1000 dilution.  
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
Lysates/proteins: 25 $\mu$ g per lane.  
Blocking buffer: 3% nonfat dry milk in TBST.  
Detection: ECL Basic Kit (RM00020).  
Exposure time: 1s.