Leader in Biomolecular Solutions for Life Science

# Reptin/RUVBL2 Rabbit pAb

Catalog No.: A1905 3 Publications



## **Basic Information**

### **Observed MW**

50kDa

### **Calculated MW**

51kDa

### Category

Mouse Monoclonal Antibody

#### **Applications**

WB,IF/ICC,IP,ELISA

### **Cross-Reactivity**

Human, Mouse, Rat

# **Background**

This gene encodes the second human homologue of the bacterial RuvB gene. Bacterial RuvB protein is a DNA helicase essential for homologous recombination and DNA double-strand break repair. Functional analysis showed that this gene product has both ATPase and DNA helicase activities. This gene is physically linked to the CGB/LHB gene cluster on chromosome 19q13.3, and is very close (55 nt) to the LHB gene, in the opposite orientation.

# **Recommended Dilutions**

**WB** 1:500 - 1:2000

**IF/ICC** 1:50 - 1:200

IP 0.5μg-4μg antibody for 200μg-400μg extracts of

whole cells

# **Immunogen Information**

 Gene ID
 Swiss Prot

 10856
 Q9Y230

### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 1-463 of human Reptin/Reptin/RUVBL2 (NP 006657.1).

### **Synonyms**

RVB2; TiH2; ECP51; TIP48; CGI-46; ECP-51; INO80J; REPTIN; TIP49B; TAP54-beta; Reptin/RUVBL2

### **Contact**

www.abclonal.com

# **Product Information**


SourceIsotypePurificationRabbitIgGAffinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

# **Validation Data**




Western blot analysis of various lysates using Reptin/RUVBL2 Rabbit pAb (A1905) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.



Immunofluorescence analysis of U2OS cells using Reptin/RUVBL2 Rabbit pAb (A1905). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunoprecipitation analysis of 200  $\mu$ g extracts of K-562 cells using 3  $\mu$ g Reptin/Reptin/RUVBL2 antibody (A1905). Western blot was performed from the immunoprecipitate using Reptin/Reptin/RUVBL2 antibody (A1905) at a dilution of 1:1000.