

A19539

Leader in Biomolecular Solutions for Life Science



## Aurora B Rabbit mAb

Catalog No.: A19539

Recombinant

5 Publications

### Basic Information

#### Observed MW

39kDa

#### Calculated MW

39kDa

#### Category

SMab Recombinant Monoclonal Antibody

#### Applications

WB, IHC-P, IF/ICC, ELISA

#### Cross-Reactivity

Human, Mouse, Rat

#### CloneNo number

ARC50905

### Background

This gene encodes a member of the aurora kinase subfamily of serine/threonine kinases. The genes encoding the other two members of this subfamily are located on chromosomes 19 and 20. These kinases participate in the regulation of alignment and segregation of chromosomes during mitosis and meiosis through association with microtubules. A pseudogene of this gene is located on chromosome 8. Alternatively spliced transcript variants have been found for this gene.

### Recommended Dilutions

WB	1:500 - 1:2000
IHC-P	1:100 - 1:500
IF/ICC	1:100 - 1:1000

### Immunogen Information

#### Gene ID

9212

#### Swiss Prot

Q96GD4

#### Immunogen

A synthetic peptide corresponding to a sequence within amino acids 1-100 of human Aurora B (NP\_004208.2).

#### Synonyms

AIK2; AIM1; ARK2; AurB; IPL1; STK5; AIM-1; ARK-2; STK-1; STK12; PPP1R48; aurkb-sv1; aurkb-sv2; Aurora B

### Contact



[www.abclonal.com](http://www.abclonal.com)

### Product Information

#### Source

Rabbit

#### Isotype

IgG

#### Purification


Affinity purification

#### Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300, 0.05% BSA, 50% glycerol, pH7.3.


## Validation Data




Western blot analysis of extracts of various cell lines, using Aurora B antibody (A19539) at 1:1000 dilution.  
 Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
 Lysates/proteins: 25µg per lane.  
 Blocking buffer: 3% nonfat dry milk in TBST.  
 Detection: ECL Basic Kit (RM00020).  
 Exposure time: 3min.



Immunohistochemistry analysis of Aurora B in paraffin-embedded mouse spleen tissue using Aurora B Rabbit mAb (A19539) at a dilution of 1:500 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of Aurora B in paraffin-embedded rat spleen tissue using Aurora B Rabbit mAb (A19539) at a dilution of 1:500 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Confocal imaging of HeLa cells using Aurora B Rabbit mAb (A19539, dilution 1:800) (Red). The cells were counterstained with Vimentin Rabbit mAb (A19607, dilution 1:100) (Green). DAPI was used for nuclear staining (blue). Objective: 60x.