Leader in Biomolecular Solutions for Life Science

# [KD Validated] DNMT1 Rabbit mAb

www.abclonal.com

ABclomal

Catalog No.: A19679

Recombinant

3 Publications

# **Basic Information**

## **Observed MW**

200kDa

### **Calculated MW**

183kDa

#### Category

Polyclonal Antibody

### **Applications**

WB,IP,ELISA

### **Cross-Reactivity**

Human, Mouse, Rat

## CloneNo number

ARC51348

# **Background**

This gene encodes an enzyme that transfers methyl groups to cytosine nucleotides of genomic DNA. This protein is the major enzyme responsible for maintaining methylation patterns following DNA replication and shows a preference for hemi-methylated DNA. Methylation of DNA is an important component of mammalian epigenetic gene regulation. Aberrant methylation patterns are found in human tumors and associated with developmental abnormalities. Variation in this gene has been associated with cerebellar ataxia, deafness, and narcolepsy, and neuropathy, hereditary sensory, type IE. Alternative splicing results in multiple transcript variants.

# **Recommended Dilutions**

**WB** 1:500 - 1:1000

IP 0.5μg-4μg antibody for 200μg-400μg extracts of

whole cells

# **Immunogen Information**

**Gene ID**Swiss Prot
1786
P26358

#### **Immunogen**

A synthetic peptide corresponding to a sequence within amino acids 1-100 of human DNMT1 (NP 001370.1).

## **Synonyms**

AIM; DNMT; MCMT; CXXC9; HSN1E; ADCADN; m.Hsal; [KD Validated] DNMT1

### **Contact**

www.abclonal.com

# **Product Information**


SourceIsotypePurificationRabbitIgGAffinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,0.05% BSA,50% glycerol,pH7.3.

# **Validation Data**



Western blot analysis of various lysates using [KD Validated] DNMT1 Rabbit mAb (A19679) at 1:1000 dilution incubated overnight at 4°C.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L)(AS014) at 1:10000 dilution.

Lysates/proteins: 25 µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020)

Exposure time: 30 s.


Western blot analysis of lysates from wild type (WT) and DNMT1 knockdown (KD) 293T cells, using [KD Validated] DNMT1 Rabbit mAb (A19679) at1:1000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% honfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).



 $Immunoprecipitation\ analysis\ of\ 300\ \mu g\ extracts\ from\ Jurkat\ cells\ using\ 3\ \mu g\ [KD\ Validated]\ DNMT1\ Rabbit$ mAb (A19679). Western stock was performed from the immunoprecipitate using DNMT1 (A19679) at a dilution of 1:10000 m. The control of the cont

