

A19879

Leader in Biomolecular Solutions for Life Science



## [KO Validated] APEH Rabbit pAb

Catalog No.: A19879 **KO** Validated

### Basic Information

#### Observed MW

81kDa

#### Calculated MW

81kDa

#### Category

Mouse Monoclonal Antibody

#### Applications

WB, ELISA

#### Cross-Reactivity

Human, Mouse, Rat

### Background

This gene encodes the enzyme acylpeptide hydrolase, which catalyzes the hydrolysis of the terminal acetylated amino acid preferentially from small acetylated peptides. The acetyl amino acid formed by this hydrolase is further processed to acetate and a free amino acid by an aminoacylase. This gene is located within the same region of chromosome 3 (3p21) as the aminoacylase gene, and deletions at this locus are also associated with a decrease in aminoacylase activity. The acylpeptide hydrolase is a homotetrameric protein of 300 kDa with each subunit consisting of 732 amino acid residues. It can play an important role in destroying oxidatively damaged proteins in living cells. Deletions of this gene locus are found in various types of carcinomas, including small cell lung carcinoma and renal cell carcinoma.

### Recommended Dilutions

WB 1:500 - 1:2000

### Immunogen Information

#### Gene ID

327

#### Swiss Prot

P13798

#### Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 1-260 of human APEH (NP\_001631.3).

#### Synonyms

APH; OPH; AARE; ACPH; D3S48E; D3F15S2; DNF15S2; EH

### Contact



[www.abclonal.com](http://www.abclonal.com)

### Product Information

#### Source

Rabbit

#### Isotype

IgG

#### Purification


Affinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal, 50% glycerol, pH 7.3.

## Validation Data



Western blot analysis of lysates from wild type (WT) and APEH knockout (KO) 293T cells, using [KO Validated] APEH Rabbit pAb (A19879) at 1:1000 dilution.  
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
Lysates/proteins: 25 $\mu$ g per lane.  
Blocking buffer: 3% nonfat dry milk in TBST.  
Detection: ECL Basic Kit (RM00020).  
Exposure time: 1s.