Leader in Biomolecular Solutions for Life Science

# TNFSF14/LIGHT Rabbit pAb

Catalog No.: A2002



### **Basic Information**

### **Observed MW**

28kDa

#### **Calculated MW**

26kDa

#### Category

Polyclonal Antibody

# **Applications**

WB, ELISA

### **Cross-Reactivity**

Human, Mouse, Rat

# **Background**

The protein encoded by this gene is a member of the tumor necrosis factor (TNF) ligand family. This protein is a ligand for TNFRSF14, which is a member of the tumor necrosis factor receptor superfamily, and which is also known as a herpesvirus entry mediator (HVEM). This protein may function as a costimulatory factor for the activation of lymphoid cells and as a deterrent to infection by herpesvirus. This protein has been shown to stimulate the proliferation of T cells, and trigger apoptosis of various tumor cells. This protein is also reported to prevent tumor necrosis factor alpha mediated apoptosis in primary hepatocyte. Two alternatively spliced transcript variant encoding distinct isoforms have been reported.

# **Recommended Dilutions**

WB

1:100 - 1:500

# Immunogen Information

Gene ID 8740 Swiss Prot 043557

043

#### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 59-176 of human TNFSF14/LIGHT (NP 003798.2).

### **Synonyms**

LTg; CD258; HVEML; LIGHT; TNFSF14/LIGHT

### **Contact**

0

www.abclonal.com

## **Product Information**

**Source** Rabbit **Isotype** IgG


**Purification**Affinity purification

## Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.

# **Validation Data**



Western blot analysis of various lysates, using TNFSF14/LIGHT antibody (A2002) at 1:500 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.