

# Versican Rabbit pAb

Catalog No.: A20278

## Basic Information

### Observed MW

Refer to figures

### Calculated MW

373kDa

### Category

Primary antibody

### Applications

ELISA, IF/ICC

### Cross-Reactivity

Human, Mouse, Rat

## Background

This gene is a member of the aggrecan/versican proteoglycan family. The protein encoded is a large chondroitin sulfate proteoglycan and is a major component of the extracellular matrix. This protein is involved in cell adhesion, proliferation, proliferation, migration and angiogenesis and plays a central role in tissue morphogenesis and maintenance. Mutations in this gene are the cause of Wagner syndrome type 1. Multiple transcript variants encoding different isoforms have been found for this gene.

## Recommended Dilutions

IF/ICC 1:50 - 1:200

## Immunogen Information

### Gene ID

1462

### Swiss Prot

P13611

### Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 1400-1586 of human Versican (NP\_004376.2).

### Synonyms

WGN; ERVR; GHAP; PG-M; WGN1; CSPG2; Versican

## Contact

 | [www.abclonal.com](http://www.abclonal.com)

## Product Information

### Source

Rabbit

### Isotype

IgG

### Purification

Affinity purification


### Storage

Store at -20°C. Avoid freeze / thaw cycles.


Buffer: PBS with 0.01% thimerosal, 50% glycerol, pH7.3.

## Validation Data

---



Immunofluorescence analysis of paraffin-embedded rat brain using Versican Rabbit pAb (A20278) at dilution of 1:100 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of paraffin-embedded mouse brain using Versican Rabbit pAb (A20278) at dilution of 1:100 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.