

A20371

Leader in Biomolecular Solutions for Life Science



CDKN2A/p16INK4a Mouse mAb

Catalog No.: A20371

Basic Information

Observed MW

17kDa

Calculated MW

8kDa/11kDa/12kDa/13kDa/16kDa/17kDa

Category

SMab Recombinant Monoclonal Antibody

Applications

WB, IHC-P, ELISA

Cross-Reactivity

Human

CloneNo number

AMC0530

Background

This gene generates several transcript variants which differ in their first exons. At least three alternatively spliced variants encoding distinct proteins have been reported, two of which encode structurally related isoforms known to function as inhibitors of CDK4 kinase. The remaining transcript includes an alternate first exon located 20 Kb upstream of the remainder of the gene; this transcript contains an alternate open reading frame (ARF) that specifies a protein which is structurally unrelated to the products of the other variants. This ARF product functions as a stabilizer of the tumor suppressor protein p53 as it can interact with, and sequester, the E3 ubiquitin-protein ligase MDM2, a protein responsible for the degradation of p53. In spite of the structural and functional differences, the CDK inhibitor isoforms and the ARF product encoded by this gene, through the regulatory roles of CDK4 and p53 in cell cycle G1 progression, share a common functionality in cell cycle G1 control. This gene is frequently mutated or deleted in a wide variety of tumors, and is known to be an important tumor suppressor gene.

Recommended Dilutions

WB	1:500 - 1:2000
IHC-P	1:50 - 1:200

Immunogen Information

Gene ID

1029

Swiss Prot

P42771/Q8N726

Immunogen

Recombinant protein of human CDKN2A/p16INK4a.

Synonyms

ARF; MLM; P14; P16; P19; CMM2; INK4; MTS1; TP16; CDK4I; CDKN2; INK4A; MTS-1; P14ARF; P19ARF; P16INK4; P16INK4A; P16-INK4A; CDKN2A/p16INK4a

Contact



www.abclonal.com

Product Information

Source

Mouse

Isotype

IgG2a, kappa

Purification


Cell culture supernatant concentrate

Storage

Store at 4°C. Avoid freeze / thaw cycles.

Tris Buffer, pH7.3-7.7, with 1% BSA and <0.1% Sodium azide.

Validation Data



Immunohistochemistry analysis of CDKN2A/p16INK4a in paraffin-embedded human cervix cancer using CDKN2A/p16INK4a Mouse mAb (A20371) at dilution of 1:50 (40x lens). Perform high pressure antigen retrieval with 10 mM Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.