

A20388

Leader in Biomolecular Solutions for Life Science



# TNKS1BP1 Rabbit pAb

Catalog No.: A20388

## Basic Information

### Observed MW

Refer to figures

### Calculated MW

182kDa

### Category

Polyclonal Antibody

### Applications

WB,IHC-P,ELISA

### Cross-Reactivity

Human,Mouse,Rat

## Background

Enables ankyrin repeat binding activity and enzyme binding activity. Involved in cellular response to ionizing radiation; double-strand break repair; and positive regulation of protein phosphorylation. Located in several cellular components, including actin cytoskeleton; adherens junction; and heterochromatin. Part of CCR4-NOT complex.

## Recommended Dilutions

|       |                |
|-------|----------------|
| WB    | 1:500 - 1:2000 |
| IHC-P | 1:50 - 1:200   |

## Immunogen Information

### Gene ID

85456

### Swiss Prot

Q9C0C2

### Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 700-1000 of human TNKS1BP1 (NP\_203754.2).

### Synonyms

TAB182; TNKS1BP1

## Contact



[www.abclonal.com](http://www.abclonal.com)

## Product Information

### Source

Rabbit

### Isotype

IgG

### Purification

Affinity purification


### Storage

Store at -20°C. Avoid freeze / thaw cycles.


Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.

## Validation Data


---



Immunohistochemistry analysis of paraffin-embedded human colon using TNKS1BP1 Rabbit pAb (A20388) at dilution of 1:200 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded mouse lung using TNKS1BP1 Rabbit pAb (A20388) at dilution of 1:200 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded rat stomach using TNKS1BP1 Rabbit pAb (A20388) at dilution of 1:200 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.