Leader in Biomolecular Solutions for Life Science

# TM4SF19 Rabbit pAb

Catalog No.: A20491 1 Publications



### **Basic Information**

### **Observed MW**

22kDa/

#### **Calculated MW**

22kDa

#### Category

Polyclonal Antibody

#### **Applications**

WB,IF/ICC,ELISA

#### **Cross-Reactivity**

Mouse

# **Background**

The protein encoded by this gene is a member of the four-transmembrane L6 superfamily. Members of this family function in various cellular processes including cell proliferation, motility, and adhesion via their interactions with integrins. In human brain tissue, this gene is expressed at high levels in the parietal lobe, occipital lobe, hippocampus, pons, white matter, corpus callosum, and cerebellum. Alternative splicing results in multiple transcript variants encoding different isoforms.

# **Recommended Dilutions**

**WB** 1:500 - 1:1000

**IF/ICC** 1:50 - 1:200

# **Immunogen Information**

**Gene ID Swiss Prot** 116211 096DZ7

#### **Immunogen**

A synthetic peptide corresponding to a sequence within amino acids 50-150 of human TM4SF19 (NP 001191827.1).

# **Synonyms**

OCTM4; TM4SF19

### **Contact**

www.abclonal.com

### **Product Information**


SourceIsotypePurificationRabbitIgGAffinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

### **Validation Data**




Western blot analysis of various lysates using TM4SF19 Rabbit pAb (A20491) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates / proteins: 25 µg per lane.

Blocking buffer: 3 % nonfat dry milk in TBST. Detection: ECL Enhanced Kit (RM00021).

Exposure time: 180s.



Immunofluorescence analysis of mouse testis using TM4SF19 Rabbit pAb (A20491) at dilution of 1:200 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.