Leader in Biomolecular Solutions for Life Science

# **DTYMK Rabbit mAb**

Catalog No.: A20921 Recombinant



## **Basic Information**

## **Observed MW**

24kDa

### **Calculated MW**

24kDa

### Category

SMab Recombinant Monoclonal Antibody

### **Applications**

WB,IF/ICC,ELISA

### **Cross-Reactivity**

Human

### CloneNo number

ARC2896

## **Background**

Enables thymidylate kinase activity. Predicted to be involved in dTDP biosynthetic process; dTTP biosynthetic process; and dUDP biosynthetic process. Predicted to act upstream of or within cellular response to growth factor stimulus and nucleotide biosynthetic process. Predicted to be located in mitochondrial intermembrane space and mitochondrial matrix. Predicted to be active in cytosol; mitochondrion; and nucleus.

## **Recommended Dilutions**

**WB** 1:100 - 1:500

IF/ICC 1:50 - 1:200

## **Immunogen Information**

**Gene ID**Swiss Prot
1841
P23919

#### **Immunogen**

A synthetic peptide corresponding to a sequence within amino acids 100-200 of human DTYMK (P23919).

## **Synonyms**

CDC8; TMPK; TYMK; CONPM; PP3731; DTYMK

## **Contact**

www.abclonal.com


## **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 0.05% BSA, 50% glycerol, pH7.3.




Western blot analysis of extracts of various cell lines, using DTYMK antibody (A20921) at 1:500 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 3s.



Immunofluorescence analysis of MCF7 cells using DTYMK Rabbit mAb (A20921) at dilution of 1:20 (40x lens). Blue: DAPI for nuclear staining.