A21058

Leader in Biomolecular Solutions for Life Science

Mineralocorticoid receptor (NR3C2) Rabbit mAb

Catalog No.: A21058 Recombinant



Basic Information

Observed MW 107kDa

Calculated MW 107kDa

Category SMab Recombinant Monoclonal Antibody

Applications WB,ELISA

Cross-Reactivity Human

CloneNo number ARC51728

Background

This gene encodes the mineralocorticoid receptor, which mediates aldosterone actions on salt and water balance within restricted target cells. The protein functions as a ligand-dependent transcription factor that binds to mineralocorticoid response elements in order to transactivate target genes. Mutations in this gene cause autosomal dominant pseudohypoaldosteronism type I, a disorder characterized by urinary salt wasting. Defects in this gene are also associated with early onset hypertension with severe exacerbation in pregnancy. Alternative splicing results in multiple transcript variants.

Recommended Dilutions

Immunogen Information

WB

1:500 - 1:1000

Gene ID 4306

Swiss Prot P08235

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 450-600 of human Mineralocorticoid receptor (NR3C2) (NP_000892.2).

Synonyms

MR; MCR; MLR; NR3C2VIT; Mineralocorticoid receptor (NR3C2)

| Contact | |
|---------|--|
|---------|--|

Product Information

www.abclonal.com

lsotype IgG Purification Affinity purification


Storage

Source

Rabbit

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,0.05% BSA,50% glycerol,pH7.3.

Validation Data



Western blot analysis of various lysates, using Mineralocorticoid receptor (NR3C2) Rabbit mAb (A21058) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (RM00021). Negative control (NC): THP-1 Exposure time: 180s.