

A21238

Leader in Biomolecular Solutions for Life Science



CD141/Thrombomodulin Rabbit mAb

Catalog No.: A21238

Recombinant

Basic Information

Observed MW

Refer to figures

Calculated MW

60kDa

Category

SMab Recombinant Monoclonal Antibody

Applications

IHC-P,IF/ICC,FC,ELISA

Cross-Reactivity

Human

CloneNo number

ARC52600

Background

The protein encoded by this intronless gene is an endothelial-specific type I membrane receptor that binds thrombin. This binding results in the activation of protein C, which degrades clotting factors Va and VIIIa and reduces the amount of thrombin generated. Mutations in this gene are a cause of thromboembolic disease, also known as inherited thrombophilia.

Recommended Dilutions

IHC-P	1:500 - 1:1000
IF/ICC	1:50 - 1:200
FC	1:100-1:500

Immunogen Information

Gene ID

7056

Swiss Prot

P07204

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 19-515 of human CD141/Thrombomodulin (NP_000352.1).

Synonyms

TM; THRM; AHUS6; BDCA3; CD141; BDCA-3; THPH12; CD141/Thrombomodulin

Contact



www.abclonal.com

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,0.05% BSA,50% glycerol,pH7.3.


Validation Data




Immunohistochemistry analysis of CD141/Thrombomodulin in paraffin-embedded human lung squamous carcinoma tissue using CD141/Thrombomodulin Rabbit mAb (A21238) at dilution of 1:1000 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.




Immunohistochemistry analysis of CD141/Thrombomodulin in paraffin-embedded human lung using CD141/Thrombomodulin Rabbit mAb (A21238) at dilution of 1:1000 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.




Immunohistochemistry analysis of CD141/Thrombomodulin in paraffin-embedded human tonsil using CD141/Thrombomodulin Rabbit mAb (A21238) at dilution of 1:1000 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.




Immunohistochemistry analysis of CD141/Thrombomodulin in paraffin-embedded human urothelial carcinoma using CD141/Thrombomodulin Rabbit mAb (A21238) at dilution of 1:1000 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.




Immunofluorescence analysis of A431 cells using CD141/Thrombomodulin Rabbit mAb (A21238) at dilution of 1:100 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of THP-1 cells using CD141/Thrombomodulin Rabbit mAb (A21238) at dilution of 1:100 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Flow cytometry: 1×10^6 Jurkat cells (negative control, left) and THP-1 cells (right) were surface-stained with CD141/Thrombomodulin Rabbit mAb (A21238, 2.5 µg/mL, orange line) or ABflo® 647 Rabbit IgG isotype control (A22070, 5 µg/mL).



Flow cytometry: 1×10^6 THP-1 cells were surface-stained with ABflo® 647 Rabbit IgG isotype control (A22070, 5 µg/mL, left) or CD141/Thrombomodulin Rabbit mAb (A21238, 2.5 µg/mL, right).

Validation Data

μl/Test, blue line), followed by Alexa Fluor®
647 conjugated goat anti-rabbit pAb
staining. Non-fluorescently stained cells
were used as blank control (red line).