Leader in Biomolecular Solutions for Life Science

# SIRT1 Rabbit pAb

Catalog No.: A21418



### **Basic Information**

#### **Observed MW**

120KD

### **Calculated MW**

82kDa

#### Category

Polyclonal Antibody

### **Applications**

WB, ELISA

#### **Cross-Reactivity**

Human, Mouse, Rat

# **Background**

This gene encodes a member of the sirtuin family of proteins, homologs to the yeast Sir2 protein. Members of the sirtuin family are characterized by a sirtuin core domain and grouped into four classes. The functions of human sirtuins have not yet been determined; however, yeast sirtuin proteins are known to regulate epigenetic gene silencing and suppress recombination of rDNA. Studies suggest that the human sirtuins may function as intracellular regulatory proteins with mono-ADP-ribosyltransferase activity. The protein encoded by this gene is included in class I of the sirtuin family. Alternative splicing results in multiple transcript variants.

# **Recommended Dilutions**

WB

1:500 - 1:1000

## **Immunogen Information**

**Gene ID** 23411

Swiss Prot 096EB6

#### **Immunogen**

A synthetic peptide corresponding to a sequence within amino acids 200-300 of human SIRT1 (NP 036370.2).

### **Synonyms**

SIR2; SIR2L1; SIR2alpha; SIRT1

### **Contact**

•

www.abclonal.com

# **Product Information**


**Source** Rabbit **Isotype** IgG **Purification**Affinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

# **Validation Data**



Western blot analysis of various lysates using SIRT1 Rabbit pAb (A21418) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 30s.