

A21962

Leader in Biomolecular Solutions for Life Science



## FLC Rabbit pAb

Catalog No.: A21962

### Basic Information

#### Observed MW

22kDa

#### Calculated MW

22kDa

#### Category

Polyclonal Antibody

#### Applications

WB, ELISA

#### Cross-Reactivity

Arabidopsis thaliana

### Background

MADS-box protein encoded by FLOWERING LOCUS C - transcription factor that functions as a repressor of floral transition and contributes to temperature compensation of the circadian clock. Expression is downregulated during cold treatment. Vernalization, FRI and the autonomous pathway all influence the state of FLC chromatin. Both maternal and paternal alleles are reset by vernalization, but their earliest activation differs in timing and location. Histone H3 trimethylation at lysine 4 and histone acetylation are associated with active FLC expression, whereas histone deacetylation and histone H3 dimethylation at lysines 9 and 27 are involved in FLC repression. Expression is also repressed by two small RNAs (30- and 24- nt) complementary to the FLC sense strand 3' to the polyA site. The small RNAs are most likely derived from an antisense transcript of FLC. Interacts with SOC1 and FT chromatin in vivo. Member of a protein complex.

### Recommended Dilutions

WB 1:100 - 1:500

### Immunogen Information

#### Gene ID

830878

#### Swiss Prot

Q957Q7

#### Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 60-196 of arabidopsis thaliana FLC (NP\_196576.1).

#### Synonyms

AGAMOUS-like 25; AGL25; FLF; FLOWERING LOCUS C; FLOWERING LOCUS F; MADS BOX PROTEIN FLOWERING LOCUS F; REDUCED STEM BRANCHING 6; RSB6; T31P16.130; T31P16\_130; FLC

### Contact



[www.abclonal.com](http://www.abclonal.com)

### Product Information

#### Source

Rabbit

#### Isotype

IgG

#### Purification


Affinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300, 50% glycerol, pH7.3.

## Validation Data



Western blot analysis of extracts of various tissues from *Arabidopsis thaliana*, using FLC Rabbit pAb (A21962) at 1:400 dilution.  
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
Lysates/proteins: 25µg per lane.  
Blocking buffer: 3% nonfat dry milk in TBST.  
Detection: ECL Basic Kit (RM00020).  
Exposure time: 0.5s.