Leader in Biomolecular Solutions for Life Science

ABflo® 488 Rabbit anti-Human CD93 mAb



Catalog No.: A23012

Basic Information

Observed MW

Calculated MW

69kDa

Category

SMab Recombinant Monoclonal Antibody

Applications

FC

Cross-Reactivity

Human

CloneNo number

ARC58668-ABf488

Conjugate

ABflo® 488. Ex:491nm. Em:516nm.

Background

The protein encoded by this gene is a cell-surface glycoprotein and type I membrane protein that was originally identified as a myeloid cell-specific marker. The encoded protein was once thought to be a receptor for C1q, but now is thought to instead be involved in intercellular adhesion and in the clearance of apoptotic cells. The intracellular cytoplasmic tail of this protein has been found to interact with moesin, a protein known to play a role in linking transmembrane proteins to the cytoskeleton and in the remodelling of the cytoskeleton.

Recommended Dilutions

FC

5 μl per 10^6 cells in 100 μl volume

Immunogen Information

Gene ID

Swiss Prot

22918 Q9NPY3

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 24-580 of human CD93 (NP_036204.2).

Synonyms

C1QR1; C1qRP; CDw93; ECSM3; MXRA4; C1qR(P); dJ737E23.1

Contact

•

www.abclonal.com

Product Information


Source Rabbit **Isotype** IgG **Purification**Affinity purification

Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.03% proclin300,0.2% BSA,pH7.3.

Validation Data












Flow cytometry:1X10^6 Jurkat cells (negative control,Left) and U-937 cells (Right) were surface-stained with ABflo® 488 Rabbit anti-Human CD93 mAb(A23012,5 µl/Test,orange line) or ABflo® 488 Rabbit lgG isotype control (A22069,5 µl/Test,blue line). Nonfluorescently stained cells was used as blank control (red line).





Flow cytometry:1X10^6 Human PBMC were surface-stained with ABflo® 488 Rabbit anti-Human CD93 mAb(A23012,5 µl/Test,orange line) or ABflo® 488 Rabbit IgG isotype control (A22069,5 µl/Test,blue line). Non-fluorescently stained cells were used as blank control (red line).



Flow cytometry:1X10^6 Human PBMC were surface-stained with ABflo® 488 Rabbit IgG isotype control (A22069,5 μ I/Test,Ieft) or ABflo® 488 Rabbit anti-Human CD93 mAb(A23012,5 μ I/Test,right).