Leader in Biomolecular Solutions for Life Science

[KD Validated] ABCG2 Rabbit pAb

Catalog No.: A23066



Basic Information

Observed MW

65-80kDa

Calculated MW

72kDa

Category

Polyclonal Antibody

Applications

WB, ELISA

Cross-Reactivity

Human, Mouse, Rat

Background

The membrane-associated protein encoded by this gene is included in the superfamily of ATP-binding cassette (ABC) transporters. ABC proteins transport various molecules across extra- and intra-cellular membranes. ABC genes are divided into seven distinct subfamilies (ABC1, MDR/TAP, MRP, ALD, OABP, GCN20, White). This protein is a member of the White subfamily. Alternatively referred to as a breast cancer resistance protein, this protein functions as a xenobiotic transporter which may play a major role in multi-drug resistance. It likely serves as a cellular defense mechanism in response to mitoxantrone and anthracycline exposure. Significant expression of this protein has been observed in the placenta, which may suggest a potential role for this molecule in placenta tissue. Multiple transcript variants encoding different isoforms have been found for this gene.

Recommended Dilutions

WB

1:500 - 1:1000

Immunogen Information

Gene ID

Swiss Prot 09UN00

Immunogen

Recombinant protein of human ABCG2.

Synonyms

MRX; MXR; ABCP; BCRP; BMDP; MXR1; ABC15; BCRP1; CD338; GOUT1; MXR-1; CDw338; CDw388; UAQTL1; EST157481; [KD Validated] ABCG2

Contact

•

www.abclonal.com

Product Information


Source Rabbit **Isotype** IgG **Purification**Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.

Validation Data



Western blot analysis of various lysates, using [KD Validated] ABCG2 Rabbit pAb (A23066) at 1:1000

dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: $25\mu g$ per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 10s.

Western blot analysis of lysates from wild type(WT) and ABCG2 knockdown (KD) HepG2 cells, using [KD Validated] ABCG2 Rabbit pAb (A23066) at 1:1000 dilution.

Secondary artifloody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

180xsates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

140 Exposure time: 10s.

