A23101

Leader in Biomolecular Solutions for Life Science

ABflo® 488 Rabbit anti-Pig CD27 mAb

Catalog No.: A23101



Basic Information

Observed MW

Calculated MW 27kDa

Category SMab Recombinant Monoclonal Antibody

Applications FC

Cross-Reactivity Pig

CloneNo number ARC57399-ABf488

Conjugate

ABflo® 488. Ex:491nm. Em:516nm.

Recommended Dilutions

FC

5 µl per 10^6 cells in 100 µl volume

Background

The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor is required for generation and long-term maintenance of T cell immunity. It binds to ligand CD70, and plays a key role in regulating B-cell activation and immunoglobulin synthesis. This receptor transduces signals that lead to the activation of NF-kappaB and MAPK8/JNK. Adaptor proteins TRAF2 and TRAF5 have been shown to mediate the signaling process of this receptor. CD27-binding protein (SIVA), a proapoptotic protein, can bind to this receptor and is thought to play an important role in the apoptosis induced by this receptor.

Immunogen Information

Gene ID 100520023 **Swiss Prot**

Immunogen

Recombinant protein of pig CD27.

Synonyms

Contact

€

www.abclonal.com

Source

Product Information

Isotype lgG


Purification Affinity purification


Storage

Rabbit

Store at 2-8°C. Avoid freeze. Buffer: PBS with 0.03% proclin300,0.2% BSA,pH7.3.

Validation Data







Flow cytometry:1X10^6 293F cells (negative control,Left) and 293F(Transfection,right) cells were surfacestained with ABflo® 488 Rabbit anti-Pig CD27 mAb(A23101,5 µl/Test,orange line) or ABflo® 488 Rabbit IgG isotype control (A22069,5 µl/Test,blue line). Nonfluorescently stained cells was used as blank control (red line). Flow cytometry:1X10^6 293F(Transfection) cells were surface-stained with ABflo® 488 Rabbit IgG isotype control (A22069,5 µl/Test,left) or ABflo® 488 Rabbit anti-Pig CD27 mAb(A23101,5 µl/Test,right).