Leader in Biomolecular Solutions for Life Science

ABflo® 647 Rabbit anti-Human Ki67 mAb



Catalog No.: A23408

Basic Information

Observed MW

Refer to figures

Calculated MW

319kDa/359kDa

Category

SMab Recombinant Monoclonal Antibody

Applications

FC (intra)

Cross-Reactivity

Human

CloneNo number

ARC57564-ABf647

Conjugate

ABflo® 647. Ex:648nm. Em:664nm.

Background

Enables protein C-terminus binding activity. Involved in regulation of chromosome segregation and regulation of mitotic nuclear division. Located in chromosome; nuclear body; and nucleolus. Colocalizes with condensed chromosome. Implicated in Crohn's disease; breast cancer; human immunodeficiency virus infectious disease; and pancreatic cancer. Biomarker of several diseases, including Barrett's esophagus; autoimmune disease of musculoskeletal system (multiple); endocrine gland cancer (multiple); gastrointestinal system cancer (multiple); and interstitial cystitis.

Recommended Dilutions

FC (intra)

5 μl per 10^6 cells in 100 μl volume

Immunogen Information

Gene ID

Swiss Prot P46013

4288

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 2245-2345 of human MKI67 (NP 002408.3).

Synonyms

KIA; MIB-; MIB-1; PPP1R105

Contact

•

www.abclonal.com

Product Information

Source Rabbit **Isotype** IgG

Purification

Affinity purification

Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.03% proclin300,0.2% BSA,pH7.3.

Validation Data









Flow cytometry:1X10^6 knockout (KO) HeLa cells(negative control,left) and Hela cells (right) were intracellularly-stained with ABflo® 647 Rabbit anti-Human Ki67 mAb(A23408,5 µl/Test,orange line) or ABflo® 647 Rabbit IgG isotype control (A22070,5 µl/Test,blue line).Nonfluorescently stained cells was used as blank control (red line).

Flow cytometry:1X10^6 Hela cells were intracellularly-stained with ABflo® 647 Rabbit IgG isotype control (A22070,5 µl/Test,left) or ABflo® 647 Rabbit anti-Human Ki67 mAb(A23408,5 µl/Test,right).