A23771

Leader in Biomolecular Solutions for Life Science

SARS-CoV-2 Spike RBD Mouse mAb

Catalog No.: A23771



Basic Information

Observed MW 30kDa

Calculated MW

Category Mouse Monoclonal Antibody

Applications WB,IF/ICC,ELISA,DB

Cross-Reactivity SARS-CoV-2

CloneNo number AMC0346R

Background

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) is an enveloped, positivesense, single-stranded RNA virus that causes coronavirus disease 2019 (COVID-19). Virus particles include the RNA genetic material and structural proteins needed for invasion of host cells. Once inside the cell the infecting RNA is used to encode structural proteins that make up virus particles, nonstructural proteins that direct virus assembly, transcription, replication and host control and accessory proteins whose function has not been determined.~ The structural proteins of SARS-CoV-2 include the envelope protein (E), spike or surface glycoprotein (S), membrane protein (M) and the nucleocapsid protein (N). The spike glycoprotein is found on the outside of the virus particle and gives coronavirus viruses their crown-like appearance. This glycoprotein mediates attachment of the virus particle and entry into the host cell. S protein is an important target for vaccine development, antibody therapies and diagnostic antigen-based tests.

Recommended Dilutions

Immunogen Information

DB	1:500 - 1:2000		
WB	1:100 - 1:500		
IF/ICC	1:50 - 1:200		

Gene ID		
43740568		

Swiss Prot

Immunogen

Recombinant fusion protein of SARS-CoV-2 CoV spike RBD.

Synonyms

spike glycoprotein; SARS-CoV-2 Spike RBD

Contact

Product Information


 www.abclonal.com

Source Mouse **lsotype** IgG2b,kappa Purification Affinity purification


Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300, 50% glycerol, pH7.3.

Validation Data



Western blot analysis of lysates from wild type (WT) and 293T cells transfected with SARS-CoV-2 Spike RBD using SARS-CoV-2 Spike RBD Mouse mAb(A23771) at 1:500 dilution. Secondary antibody:HRP Goat Anti-Mouse IgG (H+L)(AS003) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 10s.



Immobilized Recombinant SARS-CoV-2 Spike RBD Protein (RP01278LQ) at 1µg/mL (100µL/well) can bind SARS-CoV-2 Spike RBD mAb (A23771) with a linear range of 0.78-50ng/mL.