

A2421

Leader in Biomolecular Solutions for Life Science



Asymmetric DiMethyl-Histone H3-R17 Rabbit pAb

Catalog No.: A2421

7 Publications

Basic Information

Observed MW

17kDa

Calculated MW

15kDa

Category

Mouse Monoclonal Antibody

Applications

WB,IF/ICC,ELISA

Cross-Reactivity

Human,Mouse,Rat,Other (Wide Range Predicted)

Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails but instead contain a palindromic termination element. This gene is found in the small histone gene cluster on chromosome 6p22-p21.3.

Recommended Dilutions

| | |
|--------|----------------|
| WB | 1:500 - 1:1000 |
| IF/ICC | 1:50 - 1:200 |

Immunogen Information

Gene ID

8290/8350

Swiss Prot

Q16695/P68431

Immunogen

A synthetic peptide corresponding to a sequence within amino acids 1-100 of human histone H3 (NP_003520.1).

Synonyms

H3/j; H3C1; H3C2; H3C3; H3C4; H3C6; H3C7; H3C8; H3FJ; H3C10; H3C11; HIST1H3J; Asymmetric DiMethyl-Histone H3-R17

Contact



www.abclonal.com

Product Information

Source

Rabbit

Isotype

IgG

Purification



Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.


Validation Data



Dot-blot analysis of all sorts of methylation peptides using Asymmetric DiMethyl-Histone H3-R17 antibody (A2421).



Immunofluorescence analysis of 293T cells using Asymmetric DiMethyl-Histone H3-R17 Rabbit pAb (A2421). Blue: DAPI for nuclear staining.



Immunofluorescence analysis of HeLa cells using Asymmetric DiMethyl-Histone H3-R17 Rabbit pAb (A2421) at dilution of 1:100. Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.