

A2493

Leader in Biomolecular Solutions for Life Science



## FOXM1 Rabbit pAb

Catalog No.: A2493

8 Publications

### Basic Information

#### Observed MW

110kDa

#### Calculated MW

84kDa

#### Category

Polyclonal Antibody

#### Applications

WB, IF/ICC, ChIP, ELISA

#### Cross-Reactivity

Human, Mouse, Rat

### Background

The protein encoded by this gene is a transcriptional activator involved in cell proliferation. The encoded protein is phosphorylated in M phase and regulates the expression of several cell cycle genes, such as cyclin B1 and cyclin D1. Several transcript variants encoding different isoforms have been found for this gene.

### Recommended Dilutions

**WB** 1:500 - 1:1000

**IF/ICC** 1:50 - 1:200

**ChIP** 5µg antibody for  
10µg-15µg of Chromatin

### Immunogen Information

#### Gene ID

2305

#### Swiss Prot

Q08050

#### Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 29-128 of human FOXM1 (NP\_068772.2).

#### Synonyms

MPP2; HFH11; HNF-3; INS-1; MPP-2; PIG29; FKHL16; FOXM1A; FOXM1B; FOXM1C; HFH-11; TRIDENT; MPHOSPH2; FOXM1

### Contact



[www.abclonal.com](http://www.abclonal.com)

### Product Information

#### Source

Rabbit

#### Isotype

IgG

#### Purification


Affinity purification

#### Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal, 50% glycerol, pH7.3.


## Validation Data



Western blot analysis of various lysates using FOXM1 Rabbit pAb (A2493) at 1:1000 dilution.  
 Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (A5014) at 1:10000 dilution.  
 Lysates/proteins: 25µg per lane.  
 Blocking buffer: 3% nonfat dry milk in TBST.  
 Detection: ECL Basic Kit (RM00020).  
 Exposure time: 90s.



Immunofluorescence analysis of paraffin-embedded Mouse colon tissue using FOXM1 Rabbit pAb(A2493) at a dilution of 1:100 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L)(A5007) at 1:500 dilution. Blue: DAPI for nuclear staining. Perform high pressure antigen retrieval with 0.01 M citrate buffer (pH 6.0) prior to IF staining.



Chromatin immunoprecipitation analysis of extracts of 293T cells, using FOXM1 antibody (A2493) and rabbit IgG. The amount of immunoprecipitated DNA was checked by quantitative PCR. Histogram was constructed by the ratios of the immunoprecipitated DNA to the input.