Leader in Biomolecular Solutions for Life Science

# **DKK3 Rabbit pAb**

Catalog No.: A2527 1 Publications



## **Basic Information**

### **Observed MW**

38kDa

### **Calculated MW**

38kDa

### Category

Polyclonal Antibody

# **Applications**

WB,IF/ICC,ELISA

### **Cross-Reactivity**

Human, Mouse, Rat

# **Background**

This gene encodes a protein that is a member of the dickkopf family. The secreted protein contains two cysteine rich regions and is involved in embryonic development through its interactions with the Wnt signaling pathway. The expression of this gene is decreased in a variety of cancer cell lines and it may function as a tumor suppressor gene. Alternative splicing results in multiple transcript variants encoding the same protein.

# **Recommended Dilutions**

**WB** 1:500 - 1:1000

**IF/ICC** 1:50 - 1:200

# **Immunogen Information**

**Gene ID**27122

Swiss Prot
Q9UBP4

### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 22-200 of human DKK3 (NP\_056965.3).

# **Synonyms**

RIG; CRRL; REIC; DKK3

### **Contact**

www.abclonal.com


# **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.




Western blot analysis of various lysates using DKK3 Rabbit pAb (A2527) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 60s.



Immunofluorescence analysis of MCF7 cells using DKK3 Rabbit pAb (A2527) at dilution of 1:50 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.