Leader in Biomolecular Solutions for Life Science

# **TRAF5** Rabbit pAb

Catalog No.: A2536 1 Publications



### **Basic Information**

### **Observed MW**

64kDa

#### **Calculated MW**

64kDa

#### Category

Polyclonal Antibody

### **Applications**

WB, ELISA

#### **Cross-Reactivity**

Human

### **Background**

The scaffold protein encoded by this gene is a member of the tumor necrosis factor receptor-associated factor (TRAF) protein family and contains a meprin and TRAF homology (MATH) domain, a RING-type zinc finger, and two TRAF-type zinc fingers. TRAF proteins are associated with, and mediate signal transduction from members of the TNF receptor superfamily. This protein is one of the components of a multiple protein complex which binds to tumor necrosis factor (TNF) receptor cytoplasmic domains and mediates TNF-induced activation. Multiple transcript variants encoding different isoforms have been found for this gene

### **Recommended Dilutions**

WB

1:500 - 1:2000

### **Immunogen Information**

Gene ID

Swiss Prot 000463

#### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 1-80 of human TRAF5 (NP 665702.1).

### **Synonyms**

RNF84; MGC:39780; TRAF5

### **Contact**

•

www.abclonal.com

### **Product Information**


**Source** Rabbit **Isotype** IgG **Purification**Affinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

## **Validation Data**



Western blot analysis of lysates from OVCAR3 cells, using TRAF5 Rabbit pAb (A2536) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 15s.