Leader in Biomolecular Solutions for Life Science

# CD134/OX40 Rabbit pAb

Catalog No.: A3646



### **Basic Information**

### **Observed MW**

48kDa

#### **Calculated MW**

29kDa

#### Category

Polyclonal Antibody

### **Applications**

WB, ELISA

#### **Cross-Reactivity**

Rat

### **Background**

The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor has been shown to activate NF-kappaB through its interaction with adaptor proteins TRAF2 and TRAF5. Knockout studies in mice suggested that this receptor promotes the expression of apoptosis inhibitors BCL2 and BCL2IL1/BCL2-XL, and thus suppresses apoptosis. The knockout studies also suggested the roles of this receptor in CD4+ T cell response, as well as in T cell-dependent B cell proliferation and differentiation.

### **Recommended Dilutions**

WB

1:500 - 1:2000

### **Immunogen Information**

Gene ID 7293 Swiss Prot P43489

### Immunogen

A synthetic peptide corresponding to a sequence within amino acids 100-200 of human CD134/OX40 (NP\_003318.1).

### **Synonyms**

OX40; ACT35; CD134; IMD16; TXGP1L; CD134/OX40

### **Contact**

•

www.abclonal.com

### **Product Information**


**Source** Rabbit **Isotype** IgG **Purification**Affinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

## **Validation Data**



Western blot analysis of lysates from rat liver, using CD134/OX40 Rabbit pAb (A3646) at 1:3000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.