

A3861

Leader in Biomolecular Solutions for Life Science



FTO Rabbit mAb

Catalog No.: A3861

Recombinant

8 Publications

Basic Information

Observed MW

60kDa

Calculated MW

58kDa

Category

SMab Recombinant Monoclonal Antibody

Applications

WB, IHC-P, IF/ICC, IP, ELISA

Cross-Reactivity

Human

CloneNo number

ARC0851

Background

This gene is a nuclear protein of the AlkB related non-haem iron and 2-oxoglutarate-dependent oxygenase superfamily but the exact physiological function of this gene is not known. Other non-heme iron enzymes function to reverse alkylated DNA and RNA damage by oxidative demethylation. Studies in mice and humans indicate a role in nervous and cardiovascular systems and a strong association with body mass index, obesity risk, and type 2 diabetes.

Recommended Dilutions

WB	1:500 - 1:1000
IHC-P	1:50 - 1:200
IF/ICC	1:50 - 1:200
IP	0.5µg-4µg antibody for 200µg-400µg extracts of whole cells

Immunogen Information

Gene ID

79068

Swiss Prot

Q9C0B1

Immunogen

A synthetic peptide corresponding to a sequence within amino acids 1-100 of human FTO (Q9C0B1).

Synonyms

GDFF; ALKBH9; BMIQ14; FTO

Contact



www.abclonal.com

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 0.05% BSA, 50% glycerol, pH7.3.


Validation Data




Western blot analysis of various lysates using FTO Rabbit mAb (A3861) at 1:1000 dilution.
 Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (A5014) at 1:10000 dilution.
 Lysates/proteins: 25µg per lane.
 Blocking buffer: 3% nonfat dry milk in TBST.
 Detection: ECL Basic Kit (RM00020).
 Exposure time: 3min.



Immunohistochemistry analysis of FTO in paraffin-embedded human bladder cancer using FTO Rabbit mAb (A3861) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Confocal imaging of U-2 OS cells using FTO Rabbit mAb (A3861, dilution 1:50) (Red). The cells were counterstained with α-Tubulin Mouse mAb (AC012, dilution 1:400) (Green). DAPI was used for nuclear staining (blue). Objective: 100x.



Immunoprecipitation analysis of 300 µg extracts of 293T cells using 3 µg FTO antibody (A3861). Western blot was performed from the immunoprecipitate using FTO antibody (A3861) at a dilution of 1:500.