Leader in Biomolecular Solutions for Life Science

[KO Validated] PPP1CC Rabbit pAb

www.abclonal.com

ABclonal

Catalog No.: A4035

KO Validated

Basic Information

Observed MW

38kDa

Calculated MW

37kDa

Category

Polyclonal Antibody

Applications

WB,IF/ICC,ELISA

Cross-Reactivity

Human, Mouse, Rat

Background

The protein encoded by this gene belongs to the protein phosphatase family, PP1 subfamily. PP1 is an ubiquitous serine/threonine phosphatase that regulates many cellular processes, including cell division. It is expressed in mammalian cells as three closely related isoforms, alpha, beta/delta and gamma, which have distinct localization patterns. This gene encodes the gamma isozyme. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.

Recommended Dilutions

1:200 - 1:2000 WB

IF/ICC 1:50 - 1:200

Immunogen Information

Gene ID Swiss Prot 5501 P36873

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 1-323 of human PPP1CC (NP 002701.1).

Synonyms

PP1C; PP-1G; PPP1G; CC

Contact

• www.abclonal.com


Product Information

Purification Source Isotype Rabbit IgG Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.



Western blot analysis of extracts of various cell lines, using PPP1CC antibody (A4035) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.

Western blot analysis of extracts from normal (control) and PPP1CC knockout (KO) HeLa cells, using PPP1CC analysis (A4035) at 1:1000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020). Exposure time: 1s.


Exposure time: 1s.

40kDa—
—PPP1CC


35kDa—
—β-actin



Immunofluorescence analysis of H9C2 cells using [KO Validated] PPP1CC Rabbit pAb (A4035) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.



Immunofluorescence analysis of L929 cells using [KO Validated] PPP1CC Rabbit pAb (A4035) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.



Immunofluorescence analysis of U2OS cells using [KO Validated] PPP1CC Rabbit pAb (A4035) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.