

A4590

Leader in Biomolecular Solutions for Life Science



MOB4 Rabbit pAb

Catalog No.: A4590

2 Publications

Basic Information

Observed MW

26kDa

Calculated MW

26kDa

Category

Mouse Monoclonal Antibody

Applications

WB, IHC-P, ELISA

Cross-Reactivity

Human, Mouse, Rat

Background

This gene was identified based on its similarity with the mouse counterpart. Studies of the mouse counterpart suggest that the expression of this gene may be regulated during oocyte maturation and preimplantation following zygotic gene activation. Alternatively spliced transcript variants encoding distinct isoforms have been observed. Naturally occurring read-through transcription occurs between this locus and the neighboring locus HSPE1.

Recommended Dilutions

WB	1:500 - 1:2000
IHC-P	1:50 - 1:100

Immunogen Information

Gene ID

25843

Swiss Prot

Q9Y3A3

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 1-225 of human MOB4 (NP_056202.2).

Synonyms

2C4D; MOB1; MOB3; PHOCN; PREI3; CGI-95; MOBKL3; MOB4

Contact



www.abclonal.com

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal, 50% glycerol, pH7.3.


Validation Data



Western blot analysis of various lysates using MOB4 Rabbit pAb (A4590) at 1:3000 dilution.
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 5s.



Immunohistochemistry analysis of MOB4 in paraffin-embedded rat ovary using MOB4 Rabbit pAb (A4590) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunohistochemistry analysis of MOB4 in paraffin-embedded human oophoroma using MOB4 Rabbit pAb (A4590) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.