Leader in Biomolecular Solutions for Life Science

# TSC1 Rabbit mAb

Catalog No.: A5121 Recombinant



## **Basic Information**

### **Observed MW**

150kDa

### **Calculated MW**

130kDa

#### Category

SMab Recombinant Monoclonal Antibody

### **Applications**

WB, ELISA

### **Cross-Reactivity**

Human

### CloneNo number

ARC1236

## **Background**

This gene is a tumor suppressor gene that encodes the growth inhibitory protein hamartin. The encoded protein interacts with and stabilizes the GTPase activating protein tuberin. This hamartin-tuberin complex negatively regulates mammalian target of rapamycin complex 1 (mTORC1) signaling which is a major regulator of anabolic cell growth. This protein also functions as a co-chaperone for Hsp90 that inhibits its ATPase activity. This protein functions as a facilitator of Hsp90-mediated folding of kinase and non-kinase clients, including TSC2 and thereby preventing their ubiquitination and proteasomal degradation. Mutations in this gene have been associated with tuberous sclerosis and lymphangioleiomyomatosis.

## **Recommended Dilutions**

WB

1:500 - 1:1000

## Immunogen Information

Gene ID

Swiss Prot Q92574

### **Immunogen**

A synthetic peptide corresponding to a sequence within amino acids 1065-1164 of human Hamartin (Q92574).

## **Synonyms**

LAM; TSC; TSC1

### **Contact**

•

www.abclonal.com

## **Product Information**


**Source** Rabbit **Isotype** IgG **Purification**Affinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 0.05% BSA, 50% glycerol, pH7.3.

# **Validation Data**



Western blot analysis of extracts of HT-1080 cells, using TSC1 Rabbit mAb (A5121) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (RM00021).

Exposure time: 3min.