

A5182

Leader in Biomolecular Solutions for Life Science



PLC gamma 2 (PLCG2) Rabbit mAb

Catalog No.: A5182

Recombinant

1 Publications

Basic Information

Observed MW

150kDa

Calculated MW

148kDa

Category

SMab Recombinant Monoclonal Antibody

Applications

WB,IF/ICC,ELISA

Cross-Reactivity

Human,Mouse,Rat

CloneNo number

ARC1176

Background

The protein encoded by this gene is a transmembrane signaling enzyme that catalyzes the conversion of 1-phosphatidyl-1D-myo-inositol 4,5-bisphosphate to 1D-myo-inositol 1,4,5-trisphosphate (IP3) and diacylglycerol (DAG) using calcium as a cofactor. IP3 and DAG are second messenger molecules important for transmitting signals from growth factor receptors and immune system receptors across the cell membrane. Mutations in this gene have been found in autoinflammation, antibody deficiency, and immune dysregulation syndrome and familial cold autoinflammatory syndrome 3.

Recommended Dilutions

WB	1:500 - 1:1000
IF/ICC	1:50 - 1:200

Immunogen Information

Gene ID

5336

Swiss Prot

P16885

Immunogen

A synthetic peptide corresponding to a sequence within amino acids 700-800 of human PLC gamma 2 (PLC gamma 2 (PLCG2)) (P16885).

Synonyms

FCAS3; APLAID; PLC-IV; PLC-gamma-2; PLC gamma 2 (PLCG2)

Contact



www.abclonal.com

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,0.05% BSA,50% glycerol,pH7.3.


Validation Data




Western blot analysis of lysates from Raji cells, using PLC gamma 2 (PLC gamma 2 (PLCG2)) Rabbit mAb (A5182) at 1:1000 dilution.
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 10s.



Immunofluorescence analysis of Raw 264.7 cells using PLC gamma 2 (PLC gamma 2 (PLCG2)) Rabbit mAb (A5182) at dilution of 1:100 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of THP-1 cells using PLC gamma 2 (PLC gamma 2 (PLCG2)) Rabbit mAb (A5182) at dilution of 1:100 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of paraffin-embedded rat spleen using PLC gamma 2 (PLC gamma 2 (PLCG2)) Rabbit mAb (A5182) at dilution of 1:100 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.