Leader in Biomolecular Solutions for Life Science

www.abclonal.com

MonoMethyl-Histone H3-K14 Rabbit pAb

Catalog No.: A5277

Basic Information

Observed MW
17kDa

Calculated MW
16kDa

Category
Primary antibody

Applications
ELISA,WB,IF/ICC

Cross-Reactivity
Human, Mouse, Rat, Other (Wide Range Predicted)

Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is located separately from the other H3 genes that are in the histone gene cluster on chromosome 6p22-p21.3.

Recommended Dilutions

WB	1:500 - 1:1000
IF/ICC	1:20 - 1:50

Immunogen Information

Gene ID
8290/8350

Swiss Prot
Q16695/P68431

Immunogen
A synthetic monomethylated peptide around K14 of human MonoMethyl-Histone H3-K14 (NP_003520.1).

Synonyms
H3t; H3.4; H3/g; H3FT; H3C16; HIST3H3; MonoMethyl-Histone H3-K14

Contact

www.abclonal.com

Product Information


Source
Rabbit

Isotype
IgG


Purification
Affinity purification

Storage
Store at -20℃. Avoid freeze / thaw cycles.
Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.


Validation Data



Western blot analysis of various lysates using MonoMethyl-Histone H3-K14 Rabbit pAb (A5277) at 1:1000 dilution.
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 90s.



Immunofluorescence analysis of NIH-3T3 cells using MonoMethyl-Histone H3-K14 pAb (A5277) at dilution of 1:100 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of U-2 OS cells using MonoMethyl-Histone H3-K14 pAb (A5277) at dilution of 1:100 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.