CD48 Rabbit pAb

Catalog No.: A5396



Basic Information

Observed MW 45kDa

Calculated MW 28kDa

Category Polyclonal Antibody

Applications WB,ELISA

Cross-Reactivity Mouse,Rat

Recommended Dilutions

Background

This gene encodes a member of the CD2 subfamily of immunoglobulin-like receptors which includes SLAM (signaling lymphocyte activation molecules) proteins. The encoded protein is found on the surface of lymphocytes and other immune cells, dendritic cells and endothelial cells, and participates in activation and differentiation pathways in these cells. The encoded protein does not have a transmembrane domain, however, but is held at the cell surface by a GPI anchor via a C-terminal domain which maybe cleaved to yield a soluble form of the receptor. Multiple transcript variants encoding different isoforms have been found for this gene.

Immunogen Information

WB	1:500 - 1:2000	Gene ID	Swiss Prot	
		962	P09326	
		Immunogen		
		Becombinant fusion protei	Decombinant fusion protain containing a conjunce corresponding to amine acid	

Recombinant fusion protein containing a sequence corresponding to amino acids 45-210 of human CD48 (NP_001769.2).

Synonyms

BCM1; BLAST; hCD48; mCD48; BLAST1; SLAMF2; MEM-102; CD48

Contact		
Ð	www.abclonal.com	


Product Information

Source Rabbit **Isotype** IgG **Purification** Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

Validation Data



Western blot analysis of extracts of various cell lines, using CD48 antibody (A5396) at 1:1000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (RM00021).

Exposure time: 5 min.