

A5662

Leader in Biomolecular Solutions for Life Science



CD13/ANPEP Rabbit pAb

Catalog No.: A5662

5 Publications

Basic Information

Observed MW

150kDa

Calculated MW

110kDa

Category

Polyclonal Antibody

Applications

WB, IHC-P, IF/ICC, ELISA

Cross-Reactivity

Human, Mouse, Rat

Background

Aminopeptidase N is located in the small-intestinal and renal microvillar membrane, and also in other plasma membranes. In the small intestine aminopeptidase N plays a role in the final digestion of peptides generated from hydrolysis of proteins by gastric and pancreatic proteases. Its function in proximal tubular epithelial cells and other cell types is less clear. The large extracellular carboxyterminal domain contains a pentapeptide consensus sequence characteristic of members of the zinc-binding metalloproteinase superfamily. Sequence comparisons with known enzymes of this class showed that CD13 and aminopeptidase N are identical. The latter enzyme was thought to be involved in the metabolism of regulatory peptides by diverse cell types, including small intestinal and renal tubular epithelial cells, macrophages, granulocytes, and synaptic membranes from the CNS. This membrane-bound zinc metalloprotease is known to serve as a receptor for the HCoV-229E alphacoronavirus as well as other non-human coronaviruses. This gene has also been shown to promote angiogenesis, tumor growth, and metastasis and defects in this gene are associated with various types of leukemia and lymphoma.

Recommended Dilutions

WB	1:500 - 1:1000
IHC-P	1:50 - 1:200
IF/ICC	1:50 - 1:100

Immunogen Information

Gene ID

290

Swiss Prot

P15144

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 668-967 of human CD13/CD13/ANPEP (NP_001141.2).

Synonyms

APN; AP-M; AP-N; CD13; LAP1; P150; PEPN; hAPN; GP150; CD13/ANPEP

Contact



www.abclonal.com

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.



Buffer: PBS with 0.05% proclin300, 50% glycerol, pH7.3.

Validation Data




Western blot analysis of extracts of various cell lines, using CD13/ANPEP antibody (A5662) at 1:1000 dilution.
 Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
 Lysates/proteins: 25µg per lane.
 Blocking buffer: 3% nonfat dry milk in TBST.
 Detection: ECL Basic Kit (RM00020).
 Exposure time: 10s.


Western blot analysis of extracts of OVCAR3 cells, using CD13/ANPEP antibody (A5662) at 1:1000 dilution.
 Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
 Lysates/proteins: 25µg per lane.
 Blocking buffer: 3% nonfat dry milk in TBST.
 Detection: ECL Basic Kit (RM00020).
 Exposure time: 180s.



Immunohistochemistry analysis of paraffin-embedded human placenta using CD13/ANPEP Rabbit pAb (A5662) at dilution of 1:50 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.




Immunohistochemistry analysis of paraffin-embedded mouse intestine using CD13/ANPEP Rabbit pAb (A5662) at dilution of 1:50 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.




Immunohistochemistry analysis of paraffin-embedded rat kidney using CD13/ANPEP Rabbit pAb (A5662) at dilution of 1:50 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.


Validation Data



Immunofluorescence analysis of NIH/3T3 cells using CD13/CD13/ANPEP Rabbit pAb (A5662) at dilution of 1:100. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of HeLa cells using CD13/CD13/ANPEP Rabbit pAb (A5662) at dilution of 1:100. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of C6 cells using CD13/CD13/ANPEP Rabbit pAb (A5662) at dilution of 1:100. Blue: DAPI for nuclear staining.