Leader in Biomolecular Solutions for Life Science

TrKC Rabbit pAb

Catalog No.: A5842



Basic Information

Observed MW

110-115kDa

Calculated MW

94kDa

Category

Polyclonal Antibody

Applications

WB,IF/ICC,ELISA

Cross-Reactivity

Human, Mouse, Rat

Background

This gene encodes a member of the neurotrophic tyrosine receptor kinase (NTRK) family. This kinase is a membrane-bound receptor that, upon neurotrophin binding, phosphorylates itself and members of the MAPK pathway. Signalling through this kinase leads to cell differentiation and may play a role in the development of proprioceptive neurons that sense body position. Mutations in this gene have been associated with medulloblastomas, secretory breast carcinomas and other cancers. Several transcript variants encoding different isoforms have been found for this gene.

Recommended Dilutions

WB 1:500 - 1:2000

IF/ICC 1:50 - 1:200

Immunogen Information

Gene IDSwiss Prot
4916
Q16288

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 40-300 of human TrKC (NP_001007157.1).

Synonyms

TRKC; GP145-TrkC; gp145(trkC); TrKC

Contact

www.abclonal.com


Product Information

SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.




Western blot analysis of extracts of various cell lines, using TrKC antibody (A5842) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.


Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.



Immunofluorescence analysis of rat brain cells using TrKC Rabbit pAb (A5842) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.



Immunofluorescence analysis of mouse brain cells using TrKC Rabbit pAb (A5842) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.