

A6011

Leader in Biomolecular Solutions for Life Science



AQR Rabbit pAb

Catalog No.: A6011

1 Publications

Basic Information

Observed MW

171kDa

Calculated MW

171kDa

Category

Polyclonal Antibody

Applications

WB, IHC-P, ELISA

Cross-Reactivity

Human, Mouse, Rat

Background

Enables 3'-5' RNA helicase activity and single-stranded RNA binding activity. Involved in mRNA splicing, via spliceosome. Located in nucleoplasm. Part of U2-type catalytic step 2 spliceosome.

Recommended Dilutions

WB	1:500 - 1:1000
IHC-P	1:50 - 1:100

Immunogen Information

Gene ID

9716

Swiss Prot

O60306

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 400-700 of human AQR (NP_055506.1).

Synonyms

IBP160; fSAP164; AQR

Contact



www.abclonal.com

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal, 50% glycerol, pH7.3.


Validation Data




Western blot analysis of various lysates using AQR Rabbit pAb (A6011) at 1:3000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 40s.



Immunohistochemistry analysis of AQR in paraffin-embedded rat spleen using AQR Rabbit pAb (A6011) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunohistochemistry analysis of AQR in paraffin-embedded human lung cancer using AQR Rabbit pAb (A6011) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunohistochemistry analysis of AQR in paraffin-embedded mouse testis using AQR Rabbit pAb (A6011) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.