Leader in Biomolecular Solutions for Life Science

# LIN28A Rabbit pAb

Catalog No.: A6034 1 Publications



## **Basic Information**

### **Observed MW**

28kDa

### **Calculated MW**

23kDa

### Category

Polyclonal Antibody

### **Applications**

WB,IHC-P,IF/ICC,ELISA

### **Cross-Reactivity**

Human, Mouse

# **Background**

This gene encodes a LIN-28 family RNA-binding protein that acts as a posttranscriptional regulator of genes involved in developmental timing and self-renewal in embryonic stem cells. The encoded protein functions through direct interaction with target mRNAs and by disrupting the maturation of certain miRNAs involved in embryonic development. This protein prevents the terminal processing of the LET7 family of microRNAs which are major regulators of cellular growth and differentiation. Aberrant expression of this gene is associated with cancer progression in multiple tissues.

# **Recommended Dilutions**

WB	1:500 - 1:1000
IHC-P	1:50 - 1:200
IF/ICC	1:50 - 1:200

# **Immunogen Information**

Gene ID	Swiss Prot
79727	Q9H9Z2

#### **Immunogen**

A synthetic peptide corresponding to a sequence within amino acids 100-209 of human LIN28A (NP\_078950.1).

### **Synonyms**

CSDD1; LIN28; LIN-28; ZCCHC1; lin-28A; LIN28A

### **Contact**

www.abclonal.com

## **Product Information**


SourceIsotypePurificationRabbitIgGAffinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.

# **Validation Data**




Western blot analysis of lysates from T47D cells, using LIN28A Rabbit pAb (A6034) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit  $\lg G$  (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 60s.



Immunohistochemistry analysis of LIN28A in paraffin-embedded Human testicular seminoma using NY-ESO-1 Rabbit pAb (A6034) at dilution of 1:200 (40x lens).Perform high pressure antigen retrieval with 10 mM Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.