Leader in Biomolecular Solutions for Life Science

# ATGL/PNPLA2 Rabbit pAb

Catalog No.: A6245 6 Publications



### **Basic Information**

# Observed MW

55kDa

### **Calculated MW**

55kDa

#### Category

Polyclonal Antibody

### **Applications**

WB,IF/ICC,ELISA

#### **Cross-Reactivity**

Human, Mouse, Rat

# **Background**

This gene encodes an enzyme which catalyzes the first step in the hydrolysis of triglycerides in adipose tissue. Mutations in this gene are associated with neutral lipid storage disease with myopathy.

# **Recommended Dilutions**

WB

**IF/ICC** 1:50 - 1:200

1:500 - 1:1000

### **Immunogen Information**

**Gene ID**Swiss Prot
57104
Q96AD5

#### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 45-257 of human PNPLA2 (NP\_065109.1).

### **Synonyms**

ATGL; TTS2; PEDF-R; FP17548; TTS-2.2; iPLA2zeta; 1110001C14Rik; ATGL/PNPLA2

### **Contact**

www.abclonal.com


### **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.




Western blot analysis of various lysates using ATGL/PNPLA2 Rabbit pAb (A6245) at 1:500 dilution. Secondary antibody:HRP Goat Anti-Rabbit  $\log$  (H+L)(AS014) at 1:10000 dilution.

Lysates / proteins: 25 µg per lane.

Blocking buffer: 3 % nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposuretime: 60s.



Immunofluorescence analysis of U2OS cells using ATGL/PNPLA2 Rabbit pAb (A6245) at dilution of 1:100. Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.