

A6380

Leader in Biomolecular Solutions for Life Science



Gastrin Releasing Peptide (GRP) Rabbit pAb

Catalog No.: A6380

Basic Information

Observed MW

16Kda

Calculated MW

16kDa

Category

Polyclonal Antibody

Applications

WB,IHC-P,IF/ICC,ELISA

Cross-Reactivity

Human,Mouse,Rat

Background

This gene encodes a member of the bombesin-like family of gastrin-releasing peptides. The encoded preproprotein is proteolytically processed to generate two peptides, gastrin-releasing peptide and neuromedin-C. These peptides regulate numerous functions of the gastrointestinal and central nervous systems, including release of gastrointestinal hormones, smooth muscle cell contraction, and epithelial cell proliferation. These peptides are also likely to play a role in human cancers of the lung, colon, stomach, pancreas, breast, and prostate. Alternative splicing results in multiple transcript variants, at least one of which encodes a preproprotein that is proteolytically processed.

Recommended Dilutions

WB	1:500 - 1:1000
IHC-P	1:50 - 1:200
IF/ICC	1:50 - 1:200

Immunogen Information

Gene ID

2922

Swiss Prot

P07492

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 1-148 of human Gastrin Releasing Peptide (GRP) (NP_002082.2).

Synonyms

BN; GRP-10; proGRP; preproGRP; Gastrin Releasing Peptide (GRP)

Contact



www.abclonal.com

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.



Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Validation Data




Western blot analysis of extracts of various cell lines, using Gastrin Releasing Peptide (GRP) antibody (A6380) at 1:1000 dilution.
 Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
 Lysates/proteins: 25µg per lane.
 Blocking buffer: 3% nonfat dry milk in TBST.
 Detection: ECL Basic Kit (RM00020).
 Exposure time: 180s.


Western blot analysis of extracts of various cell lines, using Gastrin Releasing Peptide (GRP) antibody (A6380) at 1:1000 dilution.
 Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
 Lysates/proteins: 25µg per lane.
 Blocking buffer: 3% nonfat dry milk in TBST.
 Detection: ECL Enhanced Kit (RM00021).
 Exposure time: 180s.



Immunohistochemistry analysis of paraffin-embedded rat brain using Gastrin Releasing Peptide (GRP) Rabbit pAb (A6380) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.




Immunohistochemistry analysis of paraffin-embedded mouse heart using Gastrin Releasing Peptide (GRP) Rabbit pAb (A6380) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Immunofluorescence analysis of NIH/3T3 cells using Gastrin Releasing Peptide (GRP) Rabbit pAb (A6380) at dilution of 1:50 (40x lens). Blue: DAPI for nuclear staining.

Validation Data



Immunofluorescence analysis of PC-12 cells using Gastrin Releasing Peptide (GRP) Rabbit pAb (A6380) at dilution of 1:50 (40x lens). Blue: DAPI for nuclear staining.