

# [KO Validated] POLE3 Rabbit pAb

Catalog No.: A6469 KO Validated 4 Publications

## **Basic Information**

#### **Observed MW**

20-25kDa

# **Calculated MW**

17kDa

#### Category

Primary antibody

### **Applications**

ELISA,WB

### **Cross-Reactivity**

Human, Mouse

## **Background**

POLE3 is a histone-fold protein that interacts with other histone-fold proteins to bind DNA in a sequence-independent manner. These histone-fold protein dimers combine within larger enzymatic complexes for DNA transcription, replication, and packaging.

# **Recommended Dilutions**

WB

1:500 - 1:2000

# **Immunogen Information**

**Gene ID** 54107

**Swiss Prot** 

Q9NRF9

### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 1-147 of human POLE3 (NP\_059139.3).

### **Synonyms**

p17; YBL1; CHRAC2; CHRAC17; CHARAC17; E3

### **Contact**

**③** 

www.abclonal.com

### **Product Information**

Source Rabbit

Isotype IgG


**Purification** 

Affinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.



Western blot analysis of lysates from wild type (WT) and POLE3 knockout (KO) 293T cells, using [KO

Validated] POLE3 Rabbit pAb (A6469) at 1:1000 dilution.


Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 30s.



Western blot analysis of various lysates using [KO Validated] POLE3 Rabbit pAb (A6469) at 1:1000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (RM00021).

Exposure time: 90s.