Leader in Biomolecular Solutions for Life Science

# **HYAL2** Rabbit pAb

Catalog No.: A6624



## **Basic Information**

### **Observed MW**

54kDa

### **Calculated MW**

54kDa

### Category

Polyclonal Antibody

### **Applications**

WB, ELISA

### **Cross-Reactivity**

Human, Mouse

# **Background**

This gene encodes a weak acid-active hyaluronidase. The encoded protein is similar in structure to other more active hyaluronidases. Hyaluronidases degrade hyaluronan, one of the major glycosaminoglycans of the extracellular matrix. Hyaluronan and fragments of hyaluronan are thought to be involved in cell proliferation, migration and differentiation. Although it was previously thought to be a lysosomal hyaluronidase that is active at a pH below 4, the encoded protein is likely a GPI-anchored cell surface protein. This hyaluronidase serves as a receptor for the oncogenic virus Jaagsiekte sheep retrovirus. The gene is one of several related genes in a region of chromosome 3p21.3 associated with tumor suppression. This gene encodes two alternatively spliced transcript variants which differ only in the 5'

# **Recommended Dilutions**

WB

1:500 - 1:2000

# **Immunogen Information**

Gene ID

Swiss Prot 012891

# Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 367-473 of human HYAL2 (NP\_149348.2).

### **Synonyms**

LUCA2; HYAL2

### **Contact**

0

www.abclonal.com

## **Product Information**

**Source** Rabbit **Isotype** IgG


**Purification**Affinity purification

## Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

# **Validation Data**



Western blot analysis of various lysates using HYAL2 Rabbit pAb (A6624) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.