Leader in Biomolecular Solutions for Life Science

# **IL11RA Rabbit pAb**

Catalog No.: A6627



### **Basic Information**

### **Observed MW**

45kDa

### **Calculated MW**

45kDa

#### Category

Polyclonal Antibody

## **Applications**

WB, ELISA

#### **Cross-Reactivity**

Human, Mouse, Rat

# **Background**

Interleukin 11 is a stromal cell-derived cytokine that belongs to a family of pleiotropic and redundant cytokines that use the gp130 transducing subunit in their high affinity receptors. This gene encodes the IL-11 receptor, which is a member of the hematopoietic cytokine receptor family. This particular receptor is very similar to ciliary neurotrophic factor, since both contain an extracellular region with a 2-domain structure composed of an immunoglobulin-like domain and a cytokine receptor-like domain. Multiple alternatively spliced transcript variants have been found for this gene.

# **Recommended Dilutions**

WB

1:500 - 1:2000

# **Immunogen Information**

Gene ID 3590 Swiss Prot Q14626

#### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 30-290 of human IL11RA (NP\_001136256.1).

### **Synonyms**

CRSDA; IL11RA

### **Contact**

•

www.abclonal.com

## **Product Information**


**Source** Rabbit **Isotype** IgG **Purification**Affinity purification

## Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

# **Validation Data**



Western blot analysis of extracts of various cell lines, using IL11RA Antibody (A6627) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.